

LAUMON'S RESOLUTION OF DRINFELD'S COMPACTIFICATION IS SMALL

ALEXANDER KUZNETSOV

Let C be a smooth projective curve of genus 0. Let $\mathcal{B}$ be the variety of complete flags in an n -dimensional vector space V . Given an $(n-1)$ -tuple α of positive integers one can consider the space $\mathcal{Q}_\alpha$ of algebraic maps of degree α from C to $\mathcal{B}$. This space has drawn much attention recently in connection with Quantum Cohomology (see e.g. [Giv], [Kon]). The space $\mathcal{Q}_\alpha$ is smooth but not compact (see e.g. [Kon]). The problem of compactification of $\mathcal{Q}_\alpha$ proved very important. One compactification $\mathcal{Q}_\alpha^K$ was constructed in loc. cit. (the space of *stable maps*). Another compactification $\mathcal{Q}_\alpha^L$ (the space of *quasiflags*), was constructed in [Lau]. However, historically the first and most economical compactification $\mathcal{Q}_\alpha^D$ (the space of *quasimaps*) was constructed by Drinfeld (early 80-s, unpublished). The latter compactification is singular, while the former ones are smooth. Drinfeld has conjectured that the natural map $\pi : \mathcal{Q}_\alpha^L \rightarrow \mathcal{Q}_\alpha^D$ is a small resolution of singularities. In the present note we prove this conjecture after the necessary recollections. In fact, the proof gives some additional information about the fibers of π . It appears that every fiber has a cell decomposition, i.e. roughly speaking, is a disjoint union of affine spaces. This permits to compute not only the stalks of IC sheaf on $\mathcal{Q}_\alpha^D$ but, moreover, the Hodge structure in these stalks. Namely, the Hodge structure is a pure Tate one, and the generating function for the IC stalks is just the Lusztig's q -analogue of Kostant's partition function (see [Lus]).

In conclusion, let us mention that the Drinfeld compactifications are defined for the space of maps into flag manifolds of arbitrary semisimple group, and we plan to study their small resolutions in a paper to follow.

I am very grateful to M. Finkelberg for very useful discussions and permanent attention. I would like to thank D. Gaitsgory who has explained to me the statement of Drinfeld's conjecture.

1. THE SPACE OF MAPS INTO FLAG VARIETY

1.1. Notations. Let G be a complex semisimple simply-connected Lie group, $H \subset B$ its Cartan and Borel subgroups, N the unipotent radical of B , Y the lattice of coroots of G (with respect to H), l the rank of Y , $I = \{i_1, i_2, \dots, i_l\}$ the set of simple coroots, R^+ the set of positive coroots, X the lattice of weights, X^+ the cone of dominant weights, $\Omega = \{\omega_1, \omega_2, \dots, \omega_l\}$ the set of fundamental weights ($\langle \omega_k, i_l \rangle = \delta_{kl}$), $\mathcal{B} = G/B$ the flag variety and C a smooth projective curve of genus 0. Recall that there are canonical isomorphisms

$$H_2(\mathcal{B}, \mathbb{Z}) \cong Y \quad H^2(\mathcal{B}, \mathbb{Z}) \cong X.$$

For $\lambda \in X$ let $\mathbf{L}_\lambda$ denote the corresponding G -equivariant line bundle on $\mathcal{B}$.

I was partially supported by AMS and CRDF-2819 grants.

The map $\varphi : C \rightarrow \mathcal{B}$ has degree $\alpha \in \mathbb{N}[I] \subset Y$ if the following equivalent conditions hold:

1. $\varphi_*([C]) = \alpha$;
2. for any $\lambda \in X$ we have $\deg(\varphi^* \mathbf{L}_\lambda) = \langle \lambda, \alpha \rangle$.

We denote by $\mathcal{Q}_\alpha$ the space of algebraic maps from C to $\mathcal{B}$ of degree α . It is known that $\mathcal{Q}_\alpha$ is smooth variety and $\dim \mathcal{Q}_\alpha = 2|\alpha| + \dim \mathcal{B}$. In this paper we compare two natural compactifications of the space $\mathcal{Q}_\alpha$, which we presently describe.

1.2. Drinfeld's compactification. The Plücker embedding of the flag variety $\mathcal{B}$ gives rise to the following interpretation of $\mathcal{Q}_\alpha$.

For any irreducible representation V_λ ($\lambda \in X^+$) of G we consider the trivial vector bundle $\mathcal{V}_\lambda = V_\lambda \otimes \mathcal{O}_C$ over C .

For any G -morphism $\psi : V_\lambda \otimes V_\mu \rightarrow V_\nu$ we denote by the same letter the induced morphism $\psi : \mathcal{V}_\lambda \otimes \mathcal{V}_\mu \rightarrow \mathcal{V}_\nu$.

Then $\mathcal{Q}_\alpha$ is the space of collections of line subbundles $\mathcal{L}_\lambda \subset \mathcal{V}_\lambda$, $\lambda \in X^+$ such that:

- a) $\deg \mathcal{L}_\lambda = -\langle \lambda, \alpha \rangle$;
- b) For any nonzero G -morphism $\psi : V_\lambda \otimes V_\mu \rightarrow V_\nu$ such that $\nu = \lambda + \mu$ we have $\psi(\mathcal{L}_\lambda \otimes \mathcal{L}_\mu) = \mathcal{L}_\nu$;
- c) For any G -morphism $\psi : V_\lambda \otimes V_\mu \rightarrow V_\nu$ such that $\nu < \lambda + \mu$ we have $\psi(\mathcal{L}_\lambda \otimes \mathcal{L}_\mu) = 0$.

Remark 1.2.1. Certainly, the property b) guarantees that in order to specify such a collection it suffices to give $\mathcal{L}_{\omega_k}$ for the set Ω of fundamental weights.

If we replace the curve C by a point, we get the Plücker description of the flag variety $\mathcal{B}$ as the space of collections of lines $L_\lambda \subset V_\lambda$ satisfying conditions of type (b) and (c) (thus $\mathcal{B}$ is embedded into $\prod_{\lambda \in X^+} \mathbb{P}(V_\lambda)$). Here, a Borel subgroup $\mathbf{B}$ in

$\mathcal{B}$ corresponds to a system of lines $(L_\lambda, \lambda \in X^+)$ if lines are the fixed points of the unipotent radical of $\mathbf{B}$, $L_\lambda = (V_\lambda)^{\mathbf{N}}$, or equivalently, if $\mathbf{N}$ is the common stabilizer for all lines $\mathbf{N} = \bigcap_{\lambda \in X^+} G_{L_\lambda}$.

The following definition in case $G = SL_2$ appeared in [Dri].

Definition 1.2.2 (V.Drinfeld). The space $\mathcal{Q}_\alpha^D$ of quasimaps of degree α from C to $\mathcal{B}$ is the space of collections of invertible subsheaves $\mathcal{L}_\lambda \subset \mathcal{V}_\lambda$, $\lambda \in X^+$ such that:

- a) $\deg \mathcal{L}_\lambda = -\langle \lambda, \alpha \rangle$;
- b) For any nonzero G -morphism $\psi : V_\lambda \otimes V_\mu \rightarrow V_\nu$ such that $\nu = \lambda + \mu$ we have $\psi(\mathcal{L}_\lambda \otimes \mathcal{L}_\mu) = \mathcal{L}_\nu$;
- c) For any G -morphism $\psi : V_\lambda \otimes V_\mu \rightarrow V_\nu$ such that $\nu < \lambda + \mu$ we have $\psi(\mathcal{L}_\lambda \otimes \mathcal{L}_\mu) = 0$.

Remark 1.2.3. Here is another version of the Definition, also due to V.Drinfeld. The principal affine space $\mathcal{A} = G/N$ is an H -torsor over $\mathcal{B}$. We consider its affine closure $\mathfrak{A}$, that is, the spectrum of the ring of functions on $\mathcal{A}$. The action of H extends to $\mathfrak{A}$ but it is not free anymore. Consider the quotient stack $\tilde{\mathcal{B}} = \mathfrak{A}/H$. The flag variety $\mathcal{B}$ is an open substack in $\tilde{\mathcal{B}}$. A map $\tilde{\phi} : C \rightarrow \tilde{\mathcal{B}}$ is nothing else than an H -torsor Φ over C along with an H -equivariant morphism $f : \Phi \rightarrow \mathfrak{A}$. The degree of this map is defined as follows.

Let $\chi_\lambda : H \rightarrow \mathbb{C}^*$ be the character of H corresponding to a weight $\lambda \in X$. Let $H_\lambda \subset H$ be the kernel of the morphism χ_λ . Consider the induced $\mathbb{C}^*$ -torsor $\Phi_\lambda = \Phi/H_\lambda$ over C . The map $\tilde{\phi}$ has degree $\alpha \in \mathbb{N}[I]$ if

$$\text{for any } \lambda \in X \quad \text{we have} \quad \deg(\Phi_\lambda) = \langle \lambda, \alpha \rangle.$$

Definition 1.2.4. The space $\mathcal{Q}_\alpha^D$ is the space of maps $\tilde{\phi} : C \rightarrow \tilde{\mathcal{B}}$ of degree α such that the generic point of C maps into $\mathcal{B} \subset \tilde{\mathcal{B}}$.

The equivalence of 1.2.2 and 1.2.4 follows immediately from the Plücker embedding of $\mathfrak{A}$ into $\prod_{\lambda \in X^+} V_\lambda$.

Proposition 1.2.5. $\mathcal{Q}_\alpha^D$ is a projective variety.

Proof. The space $\mathcal{Q}_\alpha^D$ is naturally embedded into the space

$$\prod_{k=1}^l \mathbb{P}(\text{Hom}(\mathcal{O}_C(-\langle \omega_k, \alpha \rangle), \mathcal{V}_{\omega_k}))$$

and is closed in it. $\square$

1.3. The stratification of the Drinfeld's compactification. In this subsection we will introduce the stratification of the space of quasimaps.

Configurations of I -colored divisors.

Let us fix $\alpha \in \mathbb{N}[I] \subset Y$, $\alpha = \sum_{k=1}^N a_k i_k$. Consider the configuration space C^α of colored effective divisors of multidegree α (the set of colors is I). The dimension of C^α is equal to the length $|\alpha| = \sum_{k=1}^N a_k$.

Multisubsets of a set S are defined as elements of some symmetric power $S^{(m)}$ and we denote the image of $(s_1, \dots, s_m) \in S^m$ by $\{\{s_1, \dots, s_m\}\}$. We denote by $\Gamma(\alpha)$ the set of all partitions of α , i.e. multisubsets $\Gamma = \{\{\gamma_1, \dots, \gamma_m\}\}$ of $\mathbb{N}[I]$ with $\sum_{r=1}^m \gamma_r = \alpha$, $\gamma_r > 0$.

For $\Gamma \in \Gamma(\alpha)$ the corresponding stratum C_Γ^α is defined as follows. It is formed by configurations which can be subdivided into m groups of points, the r -th group containing γ_r points; all the points in one group equal to each other, the different groups being disjoint. For example, the main diagonal in C^α is the closed stratum given by partition $\alpha = \alpha$, while the complement to all diagonals in C^α is the open stratum given by partition

$$\alpha = \sum_{k=1}^N \underbrace{(i_k + i_k + \dots + i_k)}_{a_k \text{ times}}$$

Evidently, $C^\alpha = \bigsqcup_{\Gamma \in \Gamma(\alpha)} C_\Gamma^\alpha$.

Normalization and defect of subsheaves.

Let F be a vector bundle on the curve C and let E be a subsheaf in F . Let $F/E = T(E) \oplus L$ be the decomposition of the quotient sheaf F/E into the sum of its torsion subsheaf and a locally free sheaf, and let $\tilde{E} = \text{Ker}(F \rightarrow L)$ be the kernel

of the natural map $F \rightarrow L$. Then $\tilde{E}$ is a vector subbundle in F which contains E and has the following universal property:

for any subbundle $\mathcal{E}' \subset F$ if $\mathcal{E}'$ contains E then $\mathcal{E}'$ contains also $\tilde{E}$.

Moreover, $\text{rank } \tilde{E} = \text{rank } E$, $\tilde{E}/E \cong T(E)$ and $c_1(\tilde{E}) = c_1(E) + \ell(T(E))$ (for any torsion sheaf on C we denote by $\ell(T)$ its length).

Definition 1.3.1. We will call $\tilde{E}$ the *normalization* of E in F and $T(E)$ the *defect* of E .

Remark 1.3.2. If $\tilde{E}$ is the normalization of E in F then $\Lambda^k(\tilde{E})$ is the normalization of $\Lambda^k E$ in $\Lambda^k F$.

For any $x \in C$ and torsion sheaf T on C we will denote by $\ell_x(T)$ the length of the localization of T in the point x .

Definition 1.3.3. For any quasimap $\varphi = (\mathcal{L}_\lambda \subset \mathcal{V}_\lambda)_{\lambda \in X^+} \in \mathcal{Q}_\alpha^D$ we define the *normalization* of φ as follows:

$$\tilde{\varphi} = (\tilde{\mathcal{L}}_\lambda \subset \mathcal{V}_\lambda),$$

and the *defect* of φ as follows:

$$\text{def}(\varphi) = (T(\mathcal{L}_\lambda))$$

(the defect of φ is a collection of torsion sheaves).

Proposition 1.3.4. For any $\varphi \in \mathcal{Q}_\alpha^D$ there exists $\beta \leq \alpha \in \mathbb{N}[I]$, partition $\Gamma = (\gamma_1, \dots, \gamma_m) \in \Gamma(\alpha - \beta)$ and a divisor $D = \sum_{r=1}^m \gamma_r x_r \in C_\Gamma^{\alpha-\beta}$ such that

$$\tilde{\varphi} \in \mathcal{Q}_\beta \quad \ell_x(\text{def}(\varphi)_\lambda) = \begin{cases} \langle \lambda, \gamma_r \rangle, & \text{if } x = x_r \\ 0, & \text{otherwise} \end{cases}$$

Proof. Clear. □

Definition 1.3.5. The pair (β, Γ) will be called the *type of degeneration* of φ . We denote by $\mathfrak{D}_{\beta, \Gamma}$ the subspace of $\mathcal{Q}_\alpha^D$ consisting of all quasimaps φ with the given type of degeneration.

Remark 1.3.6. Note that $\mathfrak{D}_{0, \emptyset} = \mathcal{Q}_\alpha$.

We have

$$\mathcal{Q}_\alpha^D = \bigsqcup_{\substack{\beta \leq \alpha \\ \Gamma \in \Gamma(\alpha - \beta)}} \mathfrak{D}_{\beta, \Gamma} \quad (1)$$

The map $d_{\beta, \Gamma} : \mathfrak{D}_{\beta, \Gamma} \rightarrow \mathcal{Q}_\beta \times C_\Gamma^{\beta-\alpha}$ which sends φ to $(\tilde{\varphi}, D)$ (see 1.3.4) is an isomorphism. The inverse map $\sigma_{\beta, \Gamma}$ can be constructed as follows. Let $\varphi = (\mathcal{L}_\lambda) \in \mathcal{Q}_\beta$. Then

$$\sigma_{\beta, \Gamma}(\varphi, D) \stackrel{\text{def}}{=} (\mathcal{L}'_\lambda) \quad \mathcal{L}'_\lambda \stackrel{\text{def}}{=} \bigcap_{r=1}^m \mathfrak{m}_{x_r}^{\langle \lambda, \gamma_r \rangle} \cdot \mathcal{L}_\lambda,$$

where $\mathfrak{m}_x$ denotes the sheaf of ideals of the point $x \in C$.

1.4. Laumon's compactification. Let V be an n -dimensional vector space. From now on we will assume that $G = SL(V)$ (in this case certainly $l = n - 1$). In this case there is the Grassmann embedding of the flag variety, namely

$$\mathcal{B} = \{(U_1, U_2, \dots, U_{n-1}) \in G_1(V) \times G_2(V) \times \dots \times G_{n-1}(V) \mid U_1 \subset U_2 \subset \dots \subset U_{n-1}\},$$

where $G_k(V)$ is the Grassmann variety of k -dimensional subspaces in V . This embedding gives rise to another interpretation of $\mathcal{Q}_\alpha$.

We will denote by $\mathcal{V}$ the trivial vector bundle $V \otimes \mathcal{O}_C$ over C . Let $\alpha = \sum_{k=1}^{n-1} a_k i_k$, where i_k is the simple coroot dual to the highest weight ω_k of representation G in $\Lambda^k V$.

Then $\mathcal{Q}_\alpha$ is the space of complete flags of vector subbundles

$$0 \subset \mathcal{E}_1 \subset \mathcal{E}_2 \subset \dots \subset \mathcal{E}_{n-1} \subset \mathcal{V} \quad \text{such that} \quad c_1(\mathcal{E}_k) = -\langle \omega_k, \alpha \rangle = -a_k.$$

Definition 1.4.1 (Laumon, [Lau, 4.2]). The space $\mathcal{Q}_\alpha^L$ of *quasiflags* of degree α is the space of complete flags of locally free subsheaves

$$0 \subset E_1 \subset E_2 \subset \dots \subset E_{n-1} \subset \mathcal{V} \quad \text{such that} \quad c_1(E_k) = -\langle \omega_k, \alpha \rangle = -a_k.$$

It is known that $\mathcal{Q}_\alpha^L$ is a smooth projective variety of dimension $2|\alpha| + \dim \mathcal{B}$ (see loc. cit., Lemma 4.2.3).

1.5. The stratification of the Laumon's compactification. There is a stratification of the space $\mathcal{Q}_\alpha^L$ similar to the above stratification of $\mathcal{Q}_\alpha^D$.

Definition 1.5.1. For any quasiflag $E_\bullet = (E_1, \dots, E_{n-1})$ we define its *normalization* as

$$\tilde{E}_\bullet = (\tilde{E}_1, \dots, \tilde{E}_{n-1}), \quad \text{where } \tilde{E}_k \text{ is the normalization of } E_k \text{ in } \mathcal{V}$$

and *defect*

$$\text{def}(E_\bullet) = (\tilde{E}_1/E_1, \dots, \tilde{E}_{n-1}/E_{n-1})$$

Thus, the defect of $E_\bullet$ is a collection of torsion sheaves.

Proposition 1.5.2. For any $E_\bullet \in \mathcal{Q}_\alpha^L$ there exist $\beta \leq \alpha \in \mathbb{N}[I]$, partition $\Gamma = (\gamma_1, \dots, \gamma_m) \in \Gamma(\alpha - \beta)$ and a divisor $D = \sum_{r=1}^m \gamma_r x_r \in C_\Gamma^{\alpha-\beta}$ such that

$$\tilde{E}_\bullet \in \mathcal{Q}_\beta, \quad \ell_x(\text{def}(E_k)) = \begin{cases} \langle \omega_k, \gamma_r \rangle, & \text{if } x = x_r \\ 0, & \text{otherwise} \end{cases}$$

Definition 1.5.3. The pair (β, Γ) will be called the *type of degeneration* of $E_\bullet$. We denote by $\mathfrak{L}_{\beta, \Gamma}$ the subspace in $\mathcal{Q}_\alpha^L$ consisting of all quasiflags $E_\bullet$ with the given type of degeneration.

Remark 1.5.4. Note that $\mathfrak{L}_{0, \emptyset} = \mathcal{Q}_\alpha$.

We have

$$\mathcal{Q}_\alpha^L = \bigsqcup_{\substack{\beta \leq \alpha \\ \Gamma \in \Gamma(\alpha - \beta)}} \mathfrak{L}_{\beta, \Gamma} \quad (2)$$

1.6. The map from $\mathcal{Q}_\alpha^L$ to $\mathcal{Q}_\alpha^D$. Consider the map $\pi : \mathcal{Q}_\alpha^L \rightarrow \mathcal{Q}_\alpha^D$ which sends a quasiflag of degree α $E_\bullet \in \mathcal{Q}_\alpha^L$ to a quasimap given by the collection $(\mathcal{L}_{\omega_k})_{k=1}^{n-1}$ (see Remark 1.2.1) where $\mathcal{L}_{\omega_k} = \Lambda^k E_k \subset \Lambda^k \mathcal{V} = \mathcal{V}_{\omega_k}$.

Proposition 1.6.1. *Let $E_\bullet$ be a quasiflag of degree α and let (β, Γ) be its type of degeneration. Then $\pi(E_\bullet)$ is a quasimap of degree α and its type of degeneration is (β, Γ) .*

Proof. Obviously we have $\deg \mathcal{L}_{\omega_k} = \deg \Lambda^k E_k = c_1(E_k) = -\langle \omega_k, \alpha \rangle$ which means that $\pi(E_\bullet) \in \mathcal{Q}_\alpha^D$. According to the Remark 1.3.2, $\tilde{\mathcal{L}}_{\omega_k} = \Lambda^k \tilde{E}_k$ (i.e. $(\mathcal{L}_{\omega_k}) \in \mathcal{Q}_\beta$), hence

$$\ell_x(\tilde{\mathcal{L}}_{\omega_k}/\mathcal{L}_{\omega_k}) = \ell_x(\tilde{E}_k/E_k). \quad (3)$$

This proves the Proposition. $\square$

Remark 1.6.2. Note that (3) implies that π preserves not only β and Γ but also D (see 1.3.4, 1.5.2).

Recall that a proper birational map $f : \mathcal{X} \rightarrow \mathcal{Y}$ is called *small* if the following condition holds: let $\mathcal{Y}_m$ be the set of all points $y \in \mathcal{Y}$ such that $\dim f^{-1}(y) \geq m$. Then for $m > 0$ we have

$$\text{codim } \mathcal{Y}_m > 2m. \quad (4)$$

Main Theorem. The map π is a small resolution of singularities.

2. THE FIBERS OF π

2.1. We fix $\mathcal{E}_\bullet \in \mathcal{Q}_\beta$, a partition $\Gamma \in \Gamma(\alpha - \beta)$, and a divisor $D \in C_\Gamma^{\alpha-\beta}$. Then $(\mathcal{E}_\bullet, D) \in \mathfrak{D}_{\beta, \Gamma}$. We define $F(\mathcal{E}_\bullet, D)$ as $\pi^{-1}(\mathcal{E}_\bullet, D)$.

Let $D = \sum_{r=1}^m \gamma_r x_r$. We define the space $\mathcal{F}(\mathcal{E}_\bullet, D)$ of commutative diagrams

$$\begin{array}{ccccccc} \mathcal{E}_1 & \longrightarrow & \mathcal{E}_2 & \longrightarrow & \dots & \longrightarrow & \mathcal{E}_{n-1} \\ \varepsilon_1 \downarrow & & \varepsilon_2 \downarrow & & & & \varepsilon_{n-1} \downarrow \\ T_1 & \xrightarrow{\tau_1} & T_2 & \xrightarrow{\tau_2} & \dots & \xrightarrow{\tau_{n-2}} & T_{n-1} \end{array}$$

such that

- a) ε_k is surjective,
- b) T_k is torsion,
- c) $\ell_x(T_k) = \begin{cases} \langle \omega_k, \gamma_r \rangle, & \text{if } x = x_r \\ 0, & \text{otherwise} \end{cases}$

Lemma 2.1.1. *We have an isomorphism*

$$F(\mathcal{E}_\bullet, D) \cong \mathcal{F}(\mathcal{E}_\bullet, D).$$

Proof. If $E_\bullet \in F(\mathcal{E}_\bullet, D)$ then by the 1.5.2 the collection $(T_1, \dots, T_{n-1}) = \text{def}(E_\bullet)$ satisfies the above conditions.

Vice versa, if the collection $(T_1, \dots, T_k)$ satisfies the above conditions, then consider

$$E_k = \text{Ker}(\mathcal{E}_k \xrightarrow{\varepsilon_k} T_k).$$

Since the square

$$\begin{array}{ccc} \mathcal{E}_k & \xrightarrow{\varepsilon_k} & T_k \\ \downarrow & & \downarrow \tau_k \\ \mathcal{E}_{k+1} & \xrightarrow{\varepsilon_{k+1}} & T_{k+1} \end{array}$$

commutes, we can extend it to the commutative diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & E_k & \longrightarrow & \mathcal{E}_k & \xrightarrow{\varepsilon_k} & T_k & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow \tau_k & & \\ 0 & \longrightarrow & E_{k+1} & \longrightarrow & \mathcal{E}_{k+1} & \xrightarrow{\varepsilon_{k+1}} & T_{k+1} & \longrightarrow & 0 \end{array}$$

The induced morphism $E_k \rightarrow E_{k+1}$ is injective because $\mathcal{E}_k \rightarrow \mathcal{E}_{k+1}$ is, and

$$\begin{aligned} c_1(E_k) &= c_1(\mathcal{E}_k) - \ell(T_k) = -\langle \omega_k, \beta \rangle - \sum_{x \in C} \ell_x(T_k) = \\ &= -\langle \omega_k, \beta \rangle - \sum_{r=1}^m \langle \omega_k, \gamma_r \rangle = -\langle \omega_k, \beta + (\alpha - \beta) \rangle = -\langle \omega_k, \alpha \rangle \end{aligned}$$

This means that $E_\bullet \in F(\mathcal{E}_\bullet, D)$. $\square$

Proposition 2.1.2. *If $D = \sum_{r=1}^m \gamma_r x_r$ is a decomposition into disjoint divisors then*

$$F(\mathcal{E}_\bullet, D) \cong \prod_{r=1}^m F(\mathcal{E}_\bullet, \gamma_r x_r). \quad (5)$$

Proof. Recall that if T is a torsion sheaf on the curve C then

$$T = \bigoplus_{x \in C} T_x,$$

where T_x is the localization of T in the point x . This remark together with Lemma 2.1.1 proves the Proposition. $\square$

The above Proposition implies, that in order to describe general fiber $F(\mathcal{E}_\bullet, D)$ it is enough to have a description of the fibers $F(\mathcal{E}_\bullet, \gamma x)$, which we will call *simple fibers*.

2.2. The stratification of a simple fiber. We will need the following obvious Lemma.

Lemma 2.2.1. *Let $\mathcal{E}$ be a vector bundle on C . Let $\mathcal{E}' \subset \mathcal{E}$ be a vector subbundle, and let $E \subset \mathcal{E}$ be a (necessarily locally free) subsheaf. Then $E' = \mathcal{E}' \cap E$ is a vector subbundle in E .*

Moreover, the commutative square

$$\begin{array}{ccc} E' & \longrightarrow & E \\ \downarrow & & \downarrow \\ \mathcal{E}' & \longrightarrow & \mathcal{E} \end{array}$$

can be extended to the commutative diagram

$$\begin{array}{ccccc}
E' & \longrightarrow & E & \longrightarrow & E/E' \\
\downarrow & & \downarrow & & \downarrow \\
\mathcal{E}' & \longrightarrow & \mathcal{E} & \longrightarrow & \mathcal{E}/\mathcal{E}' \\
\downarrow & & \downarrow & & \downarrow \\
\mathcal{E}'/E' & \longrightarrow & \mathcal{E}/E & \longrightarrow & \frac{\mathcal{E}/E}{\mathcal{E}'/E'} \cong \frac{\mathcal{E}/\mathcal{E}'}{E/E'}
\end{array}$$

in which both the rows and the columns form the short exact sequences.

The sheaf in the lower-right corner of the diagram will be called *cointersection* of E and $\mathcal{E}'$ inside $\mathcal{E}$ and denoted by $\nabla_{\mathcal{E}}(E, \mathcal{E}')$.

Let

$$\gamma = \sum_{k=1}^{n-1} c_k i_k.$$

For every $E_{\bullet} \in F(\mathcal{E}_{\bullet}, \gamma x)$ we define

$$\mu_{pq}(E_{\bullet}) \stackrel{\text{def}}{=} l \left(\frac{\mathcal{E}_q}{E_p \cap \mathcal{E}_q} \right) \quad (1 \leq q \leq p \leq n-1), \quad (6)$$

$$\nu_{pq}(E_{\bullet}) = \begin{cases} \mu_{pq}(E_{\bullet}) - \mu_{p+1,q}(E_{\bullet}), & \text{if } 1 \leq q \leq p < n-1 \\ \mu_{pq}(E_{\bullet}), & \text{if } 1 \leq q \leq p = n-1 \end{cases} \quad (7)$$

$$\kappa_{pq}(E_{\bullet}) = \begin{cases} \nu_{pq}(E_{\bullet}) - \nu_{p,q-1}(E_{\bullet}), & \text{if } 1 < q \leq p \leq n-1 \\ \nu_{pq}(E_{\bullet}), & \text{if } 1 = q \leq p \leq n-1 \end{cases} \quad (8)$$

Remark 2.2.2. The transformations (7) and (8) are invertible, so the numbers μ_{pq} can be uniquely reconstructed from ν_{pq} or κ_{pq} . Namely,

$$\nu_{pq} = \sum_{r=1}^q \kappa_{pr}; \quad \mu_{pq} = \sum_{s=p}^{n-1} \nu_{sq} = \sum_{r \leq q \leq p \leq s} \kappa_{sr}. \quad (9)$$

Lemma 2.2.3. *We have*

$$\nu_{pq}(E_{\bullet}) = l \left(\frac{\mathcal{E}_q \cap E_{p+1}}{\mathcal{E}_q \cap E_p} \right). \quad (10)$$

$$\kappa_{pq}(E_{\bullet}) = l \left(\nabla_{\mathcal{E}_q \cap E_{p+1}}(\mathcal{E}_q \cap E_p, \mathcal{E}_{q-1} \cap E_{p+1}) \right). \quad (11)$$

Proof. The commutative diagram with exact rows

$$\begin{array}{ccccccc}
0 & \longrightarrow & \mathcal{E}_q \cap E_p & \longrightarrow & \mathcal{E}_q & \longrightarrow & \frac{\mathcal{E}_q}{\mathcal{E}_q \cap E_p} \longrightarrow 0 \\
& & \downarrow & & \parallel & & \downarrow \\
0 & \longrightarrow & \mathcal{E}_q \cap E_{p+1} & \longrightarrow & \mathcal{E}_q & \longrightarrow & \frac{\mathcal{E}_q}{\mathcal{E}_q \cap E_{p+1}} \longrightarrow 0
\end{array}$$

implies (10). In order to prove (11) note that

$$\mathcal{E}_{q-1} \cap E_p = (\mathcal{E}_q \cap E_p) \cap (\mathcal{E}_{q-1} \cap E_{p+1})$$

and apply Lemma 2.2.1 and (10). $\square$

Corollary 2.2.4. *Numbers μ_{pq} , ν_{pq} and κ_{pq} satisfy the following inequalities:*

$$0 \leq \kappa_{pq} \quad (12)$$

$$0 \leq \nu_{p1} \leq \nu_{p2} \leq \cdots \leq \nu_{pp} \quad (13)$$

$$0 \leq \mu_{n-1,q} \leq \mu_{n-2,q} \leq \cdots \leq \mu_{qq} = c_q. \quad (14)$$

Proof. See (11),(8),(7) and compare the definition of μ_{qq} with 2.1.1. $\square$

We will denote by $[p, q]$ the positive coroot

$$[p, q] \stackrel{\text{def}}{=} \sum_{k=q}^p i_k \in R^+ \quad (15)$$

Lemma 2.2.5. *For any $E_\bullet \in F(\mathcal{E}_\bullet, \gamma x)$ we have*

$$\sum_{1 \leq q \leq p \leq n-1} \kappa_{pq}(E_\bullet)[p, q] = \gamma.$$

Proof. Applying (8), (15) and (9) we get

$$\begin{aligned} \sum_{1 \leq q \leq p \leq n-1} \kappa_{pq}[p, q] &= \sum_{1 \leq q \leq p \leq n-1} (\nu_{pq} - \nu_{p,q-1})[p, q] = \\ &= \sum_{1 \leq q \leq p \leq n-1} \nu_{pq}([p, q] - [p, q+1]) = \sum_{1 \leq q \leq p \leq n-1} \nu_{pq} i_q = \sum_{q=1}^{n-1} \left(\sum_{p=q}^{n-1} \nu_{pq} \right) i_q = \sum_{q=1}^{n-1} \mu_{qq} i_q. \end{aligned}$$

Now Lemma follows from (14). $\square$

Let $\mathfrak{K}(\gamma)$ be the set of all partitions of $\gamma \in \mathbb{N}[I]$ into the sum of positive coroots: $\gamma = \sum_{s=1}^t \delta_s$, where $\delta_s \in R^+$ (note that $\mathfrak{K}(\gamma) \neq \Gamma(\gamma)$). In other words, since every positive coroot for $G = SL(V)$ is equal to $[p, q]$ for some p, q ,

$$\mathfrak{K}(\gamma) = \{(\kappa_{pq})_{1 \leq q \leq p \leq n-1} \mid \kappa_{pq} \geq 0 \text{ and } \sum_{1 \leq q \leq p \leq n-1} \kappa_{pq}[p, q] = \gamma\}.$$

Let $\mathfrak{M}(\gamma)$ denote the set of all collections (μ_{pq}) which can be produced from some $(\kappa_{pq}) \in \mathfrak{K}(\gamma)$ as in (9).

The Lemma 2.2.5 implies that for any $E_\bullet \in F(\mathcal{E}_\bullet, \gamma x)$ we have $(\mu_{pq}(E_\bullet)) \in \mathfrak{M}(\gamma)$. Define the stratum $\mathfrak{S}((\mu_{pq})_{1 \leq q \leq p \leq n-1}, (\mathcal{E}_k)_{k=1}^{n-1})$ as follows:

$$\mathfrak{S}((\mu_{pq})_{1 \leq q \leq p \leq n-1}, (\mathcal{E}_k)_{k=1}^{n-1}) = \{E_\bullet \in F(\mathcal{E}_\bullet, \gamma x) \mid \mu_{pq}(E_\bullet) = \mu_{pq}\}.$$

To unburden the notations in the cases when it is clear which flag $\mathcal{E}_\bullet$ is used we will write just $\mathfrak{S}_\mu$. We have obviously

$$F(\mathcal{E}_\bullet, \gamma x) = \bigsqcup_{\mu \in \mathfrak{M}(\gamma)} \mathfrak{S}_\mu. \quad (16)$$

Remark 2.2.6. We will also use the similar varieties $\mathfrak{S}((\mu_{pq})_{1 \leq q \leq p \leq N}, (\mathcal{E}_k)_{k=1}^N)$ that can be defined in the same way for any short flag $(\mathcal{E}_k)_{k=1}^N$ (that is the flag of subbundles $\mathcal{E}_1 \subset \cdots \subset \mathcal{E}_N \subset \mathcal{V}$ with $\text{rank } \mathcal{E}_k = k$).

2.3. The strata $\mathfrak{S}_\mu$. In order to study $\mathfrak{S}_\mu$ we will introduce some more varieties.

For every $1 \leq N \leq n-1$, a short flag of subbundles $(\mathcal{E}_k)_{k=1}^N$ (see Remark 2.2.6) and a collection of numbers $(\nu_k)_{k=1}^N$ such that $0 \leq \nu_1 \leq \dots \leq \nu_N$, we define the space $\mathfrak{T}((\nu_k)_{k=1}^N, (\mathcal{E}_k)_{k=1}^N)$ as follows:

$$\mathfrak{T}((\nu_k)_{k=1}^N, (\mathcal{E}_k)_{k=1}^N) = \{E \subset \mathcal{E}_N \mid \text{rank}(E) = N \quad \text{and} \quad l\left(\frac{\mathcal{E}_k}{\mathcal{E}_k \cap E}\right) = \nu_k\} \quad (17)$$

We define *pseudoaffine* spaces by induction in dimension. First, the affine line $\mathbb{A}^1$ is a pseudoaffine space. Now a space A is called pseudoaffine if it admits a fibration $A \rightarrow B$ with pseudoaffine fibers and pseudoaffine B .

Theorem 2.3.1. *The space $\mathfrak{T}((\nu_k)_{k=1}^N, (\mathcal{E}_k)_{k=1}^N)$ is pseudoaffine of dimension $\sum_{k=1}^{N-1} \nu_k$.*

Proof. We use induction in N . The case $N = 1$ is trivial. There is only one subsheaf E in the line bundle $\mathcal{E}_1$ with $\ell(\mathcal{E}_1/E) = \nu_1$, namely $E = \mathfrak{m}_x^{\nu_1} \cdot \mathcal{E}_1$. This means that $\mathfrak{T}(\nu_1, \mathcal{E}_1)$ is a point and the base of induction follows.

If $N > 1$ then consider the map

$$\tau : \mathfrak{T}((\nu_k)_{k=1}^N, (\mathcal{E}_k)_{k=1}^N) \rightarrow \mathfrak{T}((\nu_k)_{k=1}^{N-1}, (\mathcal{E}_k)_{k=1}^{N-1}),$$

which sends $E \in \mathfrak{T}((\nu_k)_{k=1}^N, (\mathcal{E}_k)_{k=1}^N)$ to $E' = E \cap \mathcal{E}_{N-1} \in \mathfrak{T}((\nu_k)_{k=1}^{N-1}, (\mathcal{E}_k)_{k=1}^{N-1})$.

Lemma 2.3.2. *Let $L = \mathfrak{m}_x^{\nu_N - \nu_{N-1}} \cdot \left(\frac{\mathcal{E}_N}{\mathcal{E}_{N-1}}\right)$. For any $E' \in \mathfrak{T}((\nu_k)_{k=1}^{N-1}, (\mathcal{E}_k)_{k=1}^{N-1})$ there is an isomorphism*

$$\tau^{-1}(E') \cong \text{Hom}(L, \mathcal{E}_{N-1}/E') \cong \mathbb{A}^{\ell(\mathcal{E}_{N-1}/E')} = \mathbb{A}^{\nu_{N-1}}.$$

Thus, the space $\mathfrak{T}((\nu_k)_{k=1}^N, (\mathcal{E}_k)_{k=1}^N)$ is affine fibration over a pseudoaffine space, hence it is pseudoaffine and its dimension is equal to

$$\dim(\mathfrak{T}((\nu_k)_{k=1}^{N-1}, (\mathcal{E}_k)_{k=1}^{N-1})) + \nu_{N-1} = \sum_{k=1}^{N-2} \nu_k + \nu_{N-1} = \sum_{k=1}^{N-1} \nu_k.$$

The Theorem is proved. $\square$

Proof of Lemma 2.3.2. Let $E \in \tau^{-1}(E')$. Since $E' = E \cap \mathcal{E}_{N-1}$ we can apply Lemma 2.2.1 which gives the following commutative diagram:

$$\begin{array}{ccccc} E' & \longrightarrow & E & \longrightarrow & L \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{E}_{N-1} & \xrightarrow{j} & \mathcal{E}_N & \xrightarrow{\psi} & \mathcal{E}_N/\mathcal{E}_{N-1} \\ \downarrow & & \downarrow & & \downarrow \\ T_{N-1} & \longrightarrow & T_N & \longrightarrow & T_N/T_{N-1} \end{array}$$

(Note that since $\mathcal{E}_N/\mathcal{E}_{N-1}$ is a line bundle and $\ell(T_N/T_{N-1}) = \ell(T_N) - \ell(T_{N-1}) = \nu_N - \nu_{N-1}$ the kernel of natural map $\mathcal{E}_N/\mathcal{E}_{N-1} \rightarrow T_N/T_{N-1}$ is isomorphic to L .) Let $\tilde{\mathcal{E}}_N = \psi^{-1}(L)$. Then E is contained in $\tilde{\mathcal{E}}_N$ and we have the following commutative

diagram:

$$\begin{array}{ccccc}
 E' & \longrightarrow & E & \longrightarrow & L \\
 \downarrow & & \downarrow & & \parallel \\
 \mathcal{E}_{N-1} & \xrightarrow{j} & \tilde{\mathcal{E}}_N & \xrightarrow{\psi} & L \\
 \downarrow & & \varepsilon \downarrow & & \\
 T_{N-1} & \xlongequal{\quad} & T_{N-1} & &
 \end{array}$$

This means that the points of $\tau^{-1}(E')$ are in one-to-one correspondence with maps $\varepsilon : \tilde{\mathcal{E}}_N \rightarrow T_{N-1}$ such that $\varepsilon \cdot j$ is equal to the canonical projection from $\mathcal{E}_{N-1}$ to T_{N-1} . Applying the functor $\text{Hom}(\bullet, T_{N-1})$ to the middle row of the above diagram we get an exact sequence:

$$0 \rightarrow \text{Hom}(L, T_{N-1}) \rightarrow \text{Hom}(\tilde{\mathcal{E}}_N, T_{N-1}) \xrightarrow{j^*} \text{Hom}(\mathcal{E}_{N-1}, T_{N-1}) \rightarrow \text{Ext}^1(L, T_{N-1}).$$

The last term in this sequence is zero because L is locally free and T_{N-1} is torsion. This means that the space of maps ε which we need to describe is a torsor over the group $\text{Hom}(L, T_{N-1})$. Hence this space can be identified with the group. Thus, we have proved that $\tau^{-1}(E') \cong \text{Hom}(L, T_{N-1})$ is an affine space.

Now,

$$\dim(\tau^{-1}(E')) = \dim \text{Hom}(L, T_{N-1}) = \dim H^0(T_{N-1}) = \ell(T_{N-1}) = \nu_{N-1}.$$

The Lemma is proved. $\square$

Theorem 2.3.3. *The space $\mathfrak{S}((\mu_{pq})_{1 \leq q \leq p \leq N}, (\mathcal{E}_k)_{k=1}^N)$ is a pseudoaffine space of dimension $\mu_{21} + \mu_{32} + \cdots + \mu_{N,N-1}$.*

Proof. We use induction in N . If $N = 1$ then $\mathfrak{S}_\mu$ is a point and the base of induction follows.

If $N > 1$ consider the map

$$\sigma : \mathfrak{S}((\mu_{pq})_{1 \leq q \leq p \leq N}, (\mathcal{E}_k)_{k=1}^N) \rightarrow \mathfrak{T}((\mu_{N,k})_{k=1}^N, (\mathcal{E}_k)_{k=1}^N),$$

which sends $(E_k)_{k=1}^N$ to $E_N \subset \mathcal{E}_N$.

Lemma 2.3.4. *Let $E \in \mathfrak{T}((\nu_{N,k})_{k=1}^N, (\mathcal{E}_k)_{k=1}^N)$. Consider $\tilde{\mathcal{E}}_k = \mathcal{E}_k \cap E$ ($1 \leq k \leq N-1$) and set $\tilde{\mu}_{pq} = \mu_{pq} - \mu_{Nq}$ ($1 \leq q \leq p \leq N-1$). Then $(\tilde{\mathcal{E}}_k)_{k=1}^{N-1}$ is a short flag of subbundles and for any $E \in \mathfrak{T}((\mu_{N,k})_{k=1}^N, (\mathcal{E}_k)_{k=1}^N)$*

$$\sigma^{-1}(E) \cong \mathfrak{S}((\tilde{\mu}_{pq})_{1 \leq q \leq p \leq N-1}, (\tilde{\mathcal{E}}_k)_{k=1}^{N-1}). \quad (18)$$

Thus $\mathfrak{S}((\mu_{pq})_{1 \leq q \leq p \leq N}, (\mathcal{E}_k)_{k=1}^N)$ is a fiber space with pseudoaffine base and fiber, therefore it is pseudoaffine.

Now, the calculation of the dimension

$$\begin{aligned}
 \dim(\mathfrak{S}((\mu_{pq})_{1 \leq q \leq p \leq N}, (\mathcal{E}_k)_{k=1}^N)) &= \sum_{k=1}^{N-1} \mu_{N,k} + \sum_{k=1}^{N-2} \tilde{\mu}_{k+1,k} = \\
 &= \sum_{k=1}^{N-1} \mu_{N,k} + \sum_{k=1}^{N-2} (\mu_{k+1,k} - \mu_{N,k}) = \mu_{N,N-1} + \sum_{k=1}^{N-2} \mu_{k+1,k} = \sum_{k=1}^{N-1} \mu_{k+1,k},
 \end{aligned}$$

finishes the proof of the Theorem. $\square$

Proof of Lemma 2.3.4. Assume that $(E_k)_{k=1}^N \in \mathfrak{S}_\mu$ and $E_N = E$. The commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{E}_q \cap E_p & \longrightarrow & \mathcal{E}_q & \longrightarrow & \frac{\mathcal{E}_q}{\mathcal{E}_q \cap E_p} \longrightarrow 0 \\ & & \downarrow & & \parallel & & \downarrow \\ 0 & \longrightarrow & \mathcal{E}_q \cap E & \longrightarrow & \mathcal{E}_q & \longrightarrow & \frac{\mathcal{E}_q}{\mathcal{E}_q \cap E} \longrightarrow 0 \end{array}$$

implies that

$$l\left(\frac{\mathcal{E}_q \cap E}{\mathcal{E}_q \cap E_p}\right) = l\left(\frac{\mathcal{E}_q}{\mathcal{E}_q \cap E_p}\right) - l\left(\frac{\mathcal{E}_q}{\mathcal{E}_q \cap E}\right) = \mu_{pq} - \mu_{Nq} = \tilde{\mu}_{pq}, \quad (19)$$

hence $(E_k)_{k=1}^{N-1} \in \mathfrak{S}_{\tilde{\mu}}$.

Vice versa, if $(E_k)_{k=1}^{N-1} \in \mathfrak{S}_{\tilde{\mu}}$ then the above commutative diagram along with (19) implies that $(E_k)_{k=1}^N \in \mathfrak{S}_\mu$, where we have put $E_N = E$. $\square$

2.4. The cohomology of the simple fiber. Now we will compute the dimension of the strata $\mathfrak{S}_\mu$ in terms of the partition κ .

Definition 2.4.1. A space $\mathcal{X}$ is called *cellular* if it admits a stratification with pseudoaffine strata.

Suppose $\mathcal{X} = \bigsqcup_{\xi \in \Xi} S_\xi$ is a pseudoaffine stratification of a cellular space $\mathcal{X}$. For a positive integer j we define $\chi(j) \stackrel{\text{def}}{=} \{\xi \in \Xi \mid \dim S_\xi = j\}$.

Lemma 2.4.2. *The Hodge structure $H^\bullet(\mathcal{X}, \mathbb{Q})$ is a direct sum of Tate structures, and $\mathbb{Q}(j)$ appears with multiplicity $\chi(j)$. In other words,*

$$H^\bullet(\mathcal{X}, \mathbb{Q}) = \bigoplus_{j \in \mathbb{N}} \mathbb{Q}(j)^{\chi(j)}.$$

Proof. Evident. $\square$

Given a Tate structure $\mathcal{H} = \bigoplus_{j \in \mathbb{N}} \mathbb{Q}(j)^{\chi(j)}$ we consider a *generating function*

$$P(\mathcal{H}, t) = \sum_{j \in \mathbb{N}} \chi(j) t^j \in \mathbb{N}[t].$$

For $\kappa \in \mathfrak{K}(\gamma)$ we define $K(\kappa) \stackrel{\text{def}}{=} \sum_{1 \leq q \leq p \leq n-1} \kappa_{pq}$ as the number of summands

in the partition κ . For $\gamma \in \mathbb{N}[I]$ the following q -analog of the Kostant's partition function was introduced in [Lus]:

$$\mathcal{K}_\gamma(t) = t^{|\gamma|} \sum_{\kappa \in \mathfrak{K}(\gamma)} t^{-K(\kappa)}. \quad (20)$$

Lemma 2.4.3. *Let $\kappa \in \mathfrak{K}(\gamma)$ and $\mu \in \mathfrak{M}(\gamma)$ be defined as in (9). Then*

$$\dim \mathfrak{S}_\mu = \sum_{k=1}^{n-2} \mu_{k+1,k} = |\gamma| - K(\kappa), \quad (21)$$

Proof. Applying (9) we get

$$\begin{aligned} \sum_{k=1}^{n-2} \mu_{k+1,k} &= \sum_{k=1}^{n-2} \left(\sum_{\substack{1 \leq q \leq k \\ k+1 \leq p \leq n-1}} \kappa_{pq} \right) = \sum_{1 \leq q \leq p \leq n-1} (p-q) \kappa_{pq} = \\ &= \sum_{1 \leq q \leq p \leq n-1} (|[p, q]| - 1) \kappa_{pq} = |\gamma| - \sum_{1 \leq q \leq p \leq n-1} \kappa_{pq} = |\gamma| - K(\kappa). \end{aligned}$$

□

Corollary 2.4.4. *For any $\gamma \in \mathbb{N}[I]$, $x \in C$, the simple fiber $F(\mathcal{E}_\bullet, \gamma x)$ is a cellular space, and the generating function of its cohomology is equal to the Lusztig–Kostant polynomial*

$$P(H^\bullet(F(\mathcal{E}_\bullet, \gamma x)), t) = \mathcal{K}_\gamma(t).$$

Proof. Apply (16), 2.3.3, 2.4.2 and 2.4.3. □

Corollary 2.4.5. *Let $D = \sum_{r=1}^m \gamma_r x_r \in C_\Gamma^{\alpha-\beta}$. The fiber $F(\mathcal{E}_\bullet, D)$ is a cellular space and*

$$P(H^\bullet(F(\mathcal{E}_\bullet, D)), t) = \mathcal{K}_\Gamma \stackrel{\text{def}}{=} \prod_{r=1}^m \mathcal{K}_{\gamma_r}(t). \quad (22)$$

Proof. Apply 2.1.2, 2.4.2 and 2.4.4. □

Lemma 2.4.6. *Let $D = \sum_{r=1}^m \gamma_r x_r$. We have*

$$\dim F(\mathcal{E}_\bullet, D) \leq \left\lfloor \sum_{r=1}^m \gamma_r \right\rfloor - m.$$

Proof. Note that for any $\kappa \in \mathfrak{K}(\gamma_r)$ we have $K(\kappa) \geq 1$, hence $\deg \mathcal{K}_{\gamma_r} \leq |\gamma_r| - 1$. Now, the Lemma follows from 2.4.5. □

Proof of Main Theorem. Consider the stratum $\mathfrak{D}_{\beta, \Gamma}$ of $\mathcal{Q}_\alpha^D$. Its dimension is $2|\beta| + \dim \mathcal{B} + m$ and codimension is $2|\alpha - \beta| - m$. The Lemma 2.4.6 implies that the dimension of the fiber of π over the stratum $\mathfrak{D}_{\beta, \Gamma}$ is less than or equal to $|\alpha - \beta| - m$, which is strictly less than the half codimension of the stratum. □

2.5. Applications. Let $\underline{\mathbb{Q}}$ denote the smooth constant Hodge irreducible module on $\mathcal{Q}_\alpha^L$ (as a constructible complex it lives in cohomological degree $-2|\alpha| - \dim \mathcal{B}$). Let IC denote the minimal extension of a smooth constant irreducible Hodge module from $\mathcal{Q}_\alpha$ to $\mathcal{Q}_\alpha^D$. It is well known that the smallness of π implies the following corollary.

Corollary 2.5.1.

$$IC = \pi_* \underline{\mathbb{Q}}.$$

Now we can compute the stalks of IC as cohomology of fibers of π : for $\varphi \in \mathcal{Q}_\alpha^D$ we have

$$IC_{(\varphi)}^\bullet = H^\bullet(\pi^{-1}(\varphi), \underline{\mathbb{Q}})$$

as graded Hodge structures.

Corollary 2.5.2 (Parity vanishing).

$$IC_{(\varphi)}^j = 0 \quad \text{if } j - \dim \mathcal{B} \text{ is odd.}$$

Proof. Use 2.4.5. □

Corollary 2.5.3. For $\varphi \in \mathfrak{D}_{\beta, \Gamma}$ we have

$$IC_{(\varphi)}^{-2|\alpha| - \dim \mathcal{B} + 2j} = \mathbb{Q}(j)^{\mathfrak{k}_{\Gamma}(j)},$$

where $\mathfrak{k}_{\Gamma}(j)$ is the coefficient of t^j in $\mathfrak{K}_{\Gamma}(t)$.

REFERENCES

- [Dri] Drinfeld V., Two-dimensional ℓ -adic representations of the fundamental group of a curve over a finite field and automorphic forms on $GL(2)$, American Journal of Mathematics, **105**, 1983, pp. 85–114.
- [Giv] Givental A., Equivariant Gromov–Witten Invariants, preprint, alg-geom/9603021.
- [Kon] Kontsevich M., Enumeration of rational curves via torus actions, In: The moduli space of curves, R. Dijkgraaf, C. Faber, G. van der Geer (Eds.), Progress in Math., vol. 129, Birkhäuser, 1995, pp. 335–368.
- [Lau] Laumon G., Faisceaux Automorphes Liés aux Séries d’Eisenstein, In: Automorphic Forms, Shimura Varieties, and L -functions, Perspect. Math., vol. 10, Academic Press, Boston, MA, 1990, pp. 227–281.
- [Lus] Lusztig G., Singularities, character formulas and a q -analog of weight multiplicities, Astérisque **101–102**, 1983, pp. 208–229.

INDEPENDENT UNIVERSITY OF MOSCOW, 13-26, VERESKOVAJA ST., MOSCOW 129329 RUSSIA
E-mail address: `sasha@ium.ips.ras.ru`